

Composition :-

- Each F.C tablet contains **Spiramycin 3 M. I.U.**

Properties :-

- **Unispira** is a macrolide antibiotic widely used as an alternative to the beta lactam agents for the treatment and prevention of infections caused by Gram positive organisms & infections caused by non-bacterial agents such as Chlamydia and Mycoplasma species.
- **Unispira** has also been used to treat protozoal infections such as Toxoplasmosis and Cryptosporidiosis.

Indications :-

- Community acquired E.N.T infections,
- Community acquired upper & lower respiratory Tract infections.
- Cutaneous infections.
- Prevention of infections caused by Meningococci.
- As an alternative to penicillin in case of penicillin hypersensitivity.
- Toxoplasmosis

Precautions:

Administer antibiotics, including spiramycin cautiously to any patient who has demonstrated some form of allergy, particularly to drugs.

The possibility of superinfection caused by overgrowth of nonsusceptible organisms should be kept in mind during prolonged or repeated therapy. If superinfection occurs, discontinue the drug and take appropriate measures.

Contra-Indications :-

- In case of known hypersensitivity to **Macrolides**.
- The levels of spiramycin attained in the CSF are much lower than those in the blood and are too low to be clinically useful. Therefore spiramycin must not be used in patients with meningitis.

Symptoms And Treatment Of Overdose:

Symptoms: No case of accidental overdosage has been reported. In oral doses over 4 g/day, abdominal discomfort, nausea or diarrhea may occur.

Treatment: No specific treatment has been proposed. Management should be symptomatic.

Side effects :-

- **Unispira** is well tolerated, rarely; some intestinal disturbance or some allergic reactions may occur in some patients.

Pregnancy & lactation :-

- **Unispira** can be safely used during pregnancy & used according to doctors 's advice in lactating females as it is excreted in breast milk.

Dose :-

- Usually 2 - 3 tablets daily according to physician's instructions.

Proper use of this medication :-

Taking on an empty stomach

Compliance with full course of therapy

Importance of taking medication on regular schedule and not missing doses

Proper dosing

Missed dose: taking as soon as possible; not taking if almost time for next dose; not doubling dose

Storage :-

- Keep out of the reach of children.
- Store away from heat and direct light.

Package:-

- Two strips, each contains 5 tablets.

Produced by ...

UNI PHARMA

EL OBOUR CITY, CAIRO , EGYPT

This medicament

- Is a product that affects your health and its consumption in contrary to instructions is dangerous for you .
- Follow strictly the doctor 's prescription , the stated method of administration and the instructions of the pharmacist who sold you this medication
- The doctor and the pharmacist are the experts in medicines , its benefits & risks .
- Do not interrupt the prescribed period of treatment by yourself .
- Do not repeat the medication without consulting your doctor
- Keep medicines out of reach of children

